



Editor's choice  
Scan to access more  
free content

# Contributing determinants of overall and wealth-related inequality in under-5 mortality in 13 African countries

Carine Van Malderen,<sup>1</sup> Herman Van Oyen,<sup>2</sup> Niko Speybroeck<sup>1</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jech-2012-202195>).

<sup>1</sup>Institute of Health and Society (IRSS), Université catholique de Louvain, Brussels, Belgium

<sup>2</sup>Unit of Public Health and Surveillance, Scientific Institute of Public Health, Brussels, Belgium

## Correspondence to

Carine Van Malderen, Institute of Health and Society (IRSS), Université catholique de Louvain, Clos Chapelle-aux-champs 30 bte B1.30.13, Woluwe-Saint-Lambert, Brussels 1200, Belgium, [carine.vanmalderen@uclouvain.be](mailto:carine.vanmalderen@uclouvain.be)

Received 23 November 2012

Revised 12 April 2013

Accepted 25 April 2013

Published Online First

23 May 2013

## ABSTRACT

**Objective** To investigate and compare the main determinants of overall inequality and wealth-related inequality in under-5 mortality in 13 African countries.

**Methods** Data from Demographic and Health Surveys conducted in 2007–2010 in African countries were used. The study assessed the contribution of determinants to (1) overall inequality in under-5 mortality measured by the Gini index and (2) wealth-related inequality in under-5 mortality measured by the concentration index. Techniques used were multivariate logistic regression and decomposition of Gini and concentration indexes.

**Results** Birth order and interval and region contributed the most to overall inequality in under-5 mortality in a majority of countries. A significant wealth-related inequality was observed in five countries: DR Congo, Egypt, Madagascar, Nigeria and Sao Tome & Principe. Overall, household wealth, father's occupation and mother's education contributed the most to this inequality, though the ranking of the most important determinants differed across countries.

**Conclusions** Assessing the contribution of determinants to overall inequality and to wealth-related inequality in under-5 mortality help in prioritising interventions aiming at improving child survival and equity.

## INTRODUCTION

The global under-5 death rate has dropped from 88 deaths/1000 live-births in 1990 to 57 in 2010.<sup>1</sup> However, despite substantial progress in reducing average under-5 mortality, disparity in under-5 mortality remains high: children from rural, poorer households and children of less-educated mothers are more likely to die before their fifth birthday. Tackling inequalities in health has become the new challenge. The Commission on Social Determinants of Health (CSDH) was initiated by WHO in 2005 to understand the social determinants of health, how they operate and how they can be changed to improve health and reduce health inequalities. One of the overarching recommendations of CSDH was to better measure and understand health inequality problems—within countries and globally—as a vital platform for action.<sup>2</sup>

The highest rates of under-5 mortality are observed in sub-Saharan Africa, where 1 in 8 children dies before age 5.<sup>1</sup> Accelerating the decline in under-5 mortality is possible by expanding interventions that target the most vulnerable children (eg, removing financial and social barriers to accessing basic services). Reaching the marginalised

subpopulations is therefore important not only from the point of view of social justice but also from a general public health perspective. For this purpose, information on the several dimensions of social and geographic stratification is needed.<sup>3</sup>

Once health inequalities have been observed in a population, a logical step towards guiding interventions aimed at reducing health inequalities is to understand the observed differences, for example, why do children from poor households have higher mortality? Specific analytical tools, such as a decomposition analysis,<sup>4</sup> are then needed in order to analyse the determinants of these health inequalities. Such tools allow an understanding of how a determinant affects inequality: through its more unequal distribution across the population (eg, illiteracy is more prevalent among poor mothers) or through its greater association with the health outcome (eg, mother's illiteracy is associated with infant mortality).<sup>4 5</sup>

Policy actions can be defined differently depending on whether the aim is to address the improvement of average health levels or whether it is to reduce health inequalities.<sup>6</sup> The aim of the study was to investigate the main determinants of overall inequality in under-5 mortality and of wealth-related inequality in under-5 mortality in 13 African countries.

## METHODS

### Study site and population

Data from the Demographic and Health Surveys (DHS) were used. All African countries with a standard DHS completed between 2007 and 2010 were included: DR Congo, Egypt, Ghana, Kenya, Lesotho, Liberia, Madagascar, Malawi, Nigeria, Sao Tome & Principe, Sierra Leone, Tanzania and Zambia. Egypt was the only non-sub-Saharan country selected, but was still included to allow a study of inequalities in a low under-5 mortality country. The analysis was restricted to children born alive within 5 years before the interview. Details on survey sampling, data collection and data processing can be found on the countries' final reports, available from the Measure DHS website.<sup>7</sup>

### Data and measures

The outcome variable was under-5 (0–59 months) death. This information was obtained from the birth history of interviewed women aged 15–49.<sup>7</sup> The determinants investigated were: skilled birth attendance (whether or not birth was attended by a doctor, a nurse or a midwife), sex of the child, birth order and interval between births (categorised

**To cite:** Van Malderen C, Van Oyen H, Speybroeck N. *J Epidemiol Community Health* 2013;**67**:667–676.

as: first birth; birth order 2–4 after a short (<24 months) birth interval; birth order 2–4 after a long ( $\geq 24$  months) birth interval; birth order  $\geq 5$  after a short-birth interval; birth order  $\geq 5$  after a long-birth interval), mother's age at birth (<20 years old or not), type of residence (urban or rural), region, parents' education level (none or primary and above), parents' occupation (categorised as: household/domestic, not working, unskilled manual; agriculture; other) and household wealth index (low: quintiles 1–2 or high: quintiles 3–5). The wealth index, computed by DHS, comprises household assets (type of flooring, water supply, sanitation facilities, electricity, persons per sleeping room, ownership of agricultural land, domestic servant and other assets). The choice of determinants and categories was based on results from previous studies assessing the determinants of under-5 mortality.<sup>8–16</sup> The categories of variables were willingly large to avoid analytical problems (multicollinearity resulting from the very small number of cases in several categories) that occurred in some countries.

### Ethical consideration

The study was based on data from the DHS programme. What is done to maintain high ethical standards is described in ref.<sup>17</sup>

### Data analysis

Datasets were transferred to RGui (R V2.14.2., The R foundation for Statistical Computing) for analysis.

The concentration index (C)<sup>18</sup> was used as a measure of socioeconomic inequality: briefly, a relative concentration curve plot on the x-axis, the cumulative proportion of children ranked by household wealth, and on the y-axis the cumulative proportion of the variable of interest (under-5 mortality or a determinant). C is defined as twice the area between the concentration curve and the diagonal (line of equality). In case of maximal equality, the concentration curve lies on the diagonal and C equals 0. If the curve is above the diagonal, C is negative and the variable of interest is more prevalent among the poorer households. The value of C ranges between -1 and 1. However, the bounds of C for a dichotomous variable depend on the mean of the variable.<sup>19</sup> Given that, a normalised C (C\*) proposed by Erreygers was computed when comparing countries.

Univariate and multivariate analyses were performed by country. Under-5 mortality was first computed across categories of determinants, and proportions were compared using  $\chi^2$  test. Then, a multivariate logistic regression model was used. Explanatory variables were: low household wealth, absence of skilled birth attendance, male sex, birth order and interval categories, mother's age at birth <20 years, rural residence, low parents' education and parents' occupational categories. Dummies were created for the categories of each variable and the analysis was performed on dummies. As father characteristics were included, analyses were restricted to children from mothers who ever married/lived with a man (above 93% of children in all other countries except in Liberia: 87% and Lesotho: 88%). Multicollinearity was checked and no variance inflation factor was greater than 10. The criterion used for statistical significance was  $\alpha=0.01$ . No selection of models was carried out as the objective was to study the contribution of all the determinants included. Analyses were weighted (weights provided with the DHS data) and adjusted for the cluster randomised sampling frame (with cluster as the primary sampling unit and household as the secondary sampling unit).

The decomposition analysis was performed on predicted (ln odds) under-5 mortality given the observed values of the determinants to satisfy the linearity assumption of the decomposition

method.<sup>20</sup> The overall inequality in predicted under-5 mortality measured by the Gini index (G)<sup>21</sup> and the wealth-related inequality in predicted under-5 mortality measured by C<sup>18</sup> were decomposed using Wagstaff's method.<sup>20</sup> The method is described elsewhere in detail.<sup>4 5 20 22</sup> G is similar to C with the socioeconomic variable being replaced by the health variable itself. In other words, the cumulative proportion of predicted under-5 mortality is used on the y-axis and the cumulative proportion of children ranked by predicted under-5 mortality is used on the x-axis. As the analysis is performed on predicted under-5 mortality values having a negative mean, the value of G ranges between 0 and -1. In addition, as an alternative method for assessing the contributions of determinants to the overall variability in under-5 mortality, the variance of predicted under-5 mortality was decomposed using Shorrocks' method.<sup>23</sup> Finally, contributions of determinants were ranked by importance and median ranks across countries were calculated.

## RESULTS

### Countries' profile: geographical inequalities in average under-5 mortality


Figure 1 maps the national and regional proportions of under-5 deaths in the 13 countries. This figure highlights the differences that exist between countries and within countries. Some countries have a uniformly distributed low (<6% in all regions in Egypt, Sao Tome & Principe) or high (>8% in all regions in Sierra Leone, Lesotho) under-5 mortality. But in other countries, some regions stand out. For instance, the South West region of Nigeria shows a relatively low figure compared with the rest of the country (table 1,  $p<0.01$ ), and the Nyanza region in Kenya shows a relatively high figure compared with the rest of the country (table 1,  $p<0.01$ ).

### Countries' average under-5 mortality and wealth-related inequality

Figure 2 compares the 13 countries in terms of wealth-related inequality in under-5 mortality, measured by C\*, and average under-5 mortality. With the exception of Tanzania, Malawi and Zambia, all countries showed a negative C\*, under-5 mortality being more concentrated among the poorer households. The measure was significant (upper limit of the CI being below 0) in five countries: DR Congo, Egypt, Madagascar, Nigeria and Sao Tome & Principe. Under-5 mortality ranged from 2.48% in Egypt to 11.14% in Nigeria. Across all countries, the median under-5 mortality was 8.25% (Interquartile range: 6.12%–10.29%). Among countries with the lowest under-5 mortality, Sao Tome & Principe had the highest level of inequality compared with Egypt and Madagascar (C\*=-0.032 (-0.052 to -0.013), -0.012 (-0.018 to -0.005) and -0.013 (-0.022 to -0.004), respectively). Among countries with the highest under-5 mortality, DR Congo and Nigeria had the highest level of inequality compared with Sierra Leone and Lesotho (C\*=-0.050 (-0.065 to -0.035), -0.057 (-0.065 to -0.050), -0.007 (-0.027 to 0.013) and -0.018 (-0.038 to 0.001), respectively).

### Wealth related-inequality in under-5 mortality: patterns

Figure 3 plots average under-5 mortality by wealth quintiles in the five countries where the concentration index was significant. The obtained patterns illustrate the form of wealth inequity: 'mass deprivation'—the entire population is affected except the richest minority, 'exclusion'—only the poorest minority is affected or 'queuing'—middle-income and upper-income groups are better-off than low-income groups.<sup>24</sup> For Nigeria, the curve can be compared with the 'queuing' pattern of inequality:



**Figure 1** Proportion of under-5 deaths (%) in the 13 countries: national and regional levels, Demographic and Health Surveys 2007–2010. Quantiles were calculated across all regions from the 13 countries.

under-5 mortality exceeds 12% in the majority of the population while children from quintile 4 and particularly quintile 5 have a much better survival. The DRCongo and Sao Tome & Principe patterns show a similar ‘queuing’ pattern except that in these two countries quintile 3 seems less affected than quintile 4. Finally, in Madagascar and Egypt, under-5 mortality follows a more linear gradient across wealth quintiles.

#### Determinants associated with under-5 mortality in the univariate and multivariate analyses

Under-5 mortality by determinants’ categories in each country is presented in table 1, and CIs are presented in an online supplementary appendix. Considering countries with significant differences across categories of determinants, under-5 mortality was higher in children from poorer households; with no skilled birth attendance; being male; with a short (<24 months) preceding birth interval; first or  $\geq 5$  in the birth order; living in rural residence; from parents with a low education level; or from parents working in the agricultural sector. In six countries, under-5 mortality also differed by region. Regression coefficients of the multivariate logistic regression are presented in table 2. When controlling for the other variables, in most countries, mortality was higher for male children; with a short preceding birth interval; or first in the birth order. The remaining determinants showed heterogeneous effects across countries.

#### Determinants’ contributions to overall inequality and to wealth-related inequality in under-5 mortality

The more a determinant is prevalent, associated with the outcome and unequally distributed, the more it contributes to inequality. The contribution of determinants to overall inequality and to wealth-related inequality in under-5 mortality is shown in table 3. The first column shows the Gini index (above) and the concentration index (below) for each country. The following columns show contributions by determinants, adding up to 100%. For instance, in Egypt, birth order and interval contributed to 35% of the overall inequality in under-5 mortality and to 33.4% to wealth-related inequality in under-5 mortality.

First, overall inequality in under-5 mortality, measured by the Gini index, was decomposed into its contributing determinants. Birth order and interval was the most important determinant in a majority of countries. But in Ghana, Kenya and Tanzania, most of the overall inequality in under-5 mortality was explained by regional differences. In Sao Tome & Principe, the determinant contributing most to the overall inequality was mother’s education. Among the remaining determinants, the contributions exceeding 10% included: household wealth (Egypt), skilled birth attendance (Lesotho), sex of the child (Lesotho, Liberia, Malawi, Sao Tome & Principe), mother’s age at birth (Lesotho, Liberia), type of residence (Nigeria, Sao Tome & Principe), father’s education (Ghana, Kenya, Liberia), mother’s occupation (Liberia) and father’s occupation (Egypt, Ghana). Skilled birth attendance appeared to contribute less to

**Table 1** Proportion of under-5 deaths (%) by determinant in the 13 countries (Demographic and Health Surveys 2007–2010)


	DRCongo 2007	Egypt 2008	Ghana 2008	Kenya 2008/2009	Lesotho 2009/2010	Liberia 2007	Madagascar 2008/2009	Malawi 2010	Nigeria 2008	Sao Tome & Principe 2008/2009	Sierra Leone 2008	Tanzania 2009/2010	Zambia 2007
N (10 <sup>3</sup> )	9.0	10.9	3.0	6.1	4.0	5.8	12.4	20.0	28.7	1.9	5.6	8.0	6.4
All	11.0	2.5	6.1	6.3	10.3	8.2	5.6	8.6	11.1	4.6	10.3	6.2	8.9
Quintile of household wealth index													
1	<b>13.5</b>	3.0	6.8	7.2	10.3	9.7	6.3	7.6	<b>13.7</b>	6.7	12.7	6.4	8.3
2	<b>13.1</b>	3.0	4.8	6.3	12.0	7.4	6.2	9.6	<b>13.0</b>	5.9	8.0	5.7	8.5
3	<b>9.7</b>	2.4	7.7	7.2	8.5	7.9	5.2	8.3	<b>11.6</b>	3.5	10.1	6.9	8.8
4	<b>11.1</b>	2.3	5.8	3.9	12.5	8.7	5.1	9.6	<b>9.4</b>	3.9	9.7	5.4	10.3
5 ( <i>richest</i> )	<b>6.4</b>	1.6	5.2	6.7	7.8	7.2	4.6	7.6	<b>6.4</b>	2.7	11.0	7.0	9.1
Skilled birth attendance													
Yes	<b>8.4</b>	2.4	5.8	6.3	9.0	7.7	4.7	8.3	<b>8.5</b>	3.4	8.5	6.0	9.0
No	<b>12.1</b>	2.9	6.1	6.1	12.4	8.2	5.9	8.9	<b>11.9</b>	2.7	10.2	5.9	8.6
Sex of the child													
Female	10.2	2.1	5.7	5.6	<b>8.3</b>	7.5	5.3	7.5	<b>10.5</b>	1.9	10.5	5.4	7.9
Male	11.9	2.8	6.5	7.0	<b>12.2</b>	8.9	5.8	<b>9.6</b>	<b>11.7</b>	7.2	10.1	7.0	9.9
Birth order and preceding birth interval													
First	<b>12.5</b>	<b>2.4</b>	<b>6.0</b>	5.8	<b>9.5</b>	9.8	<b>5.4</b>	<b>9.6</b>	<b>10.4</b>	<b>2.4</b>	<b>12.1</b>	7.0	<b>10.2</b>
2–4 short	<b>14.0</b>	<b>4.4</b>	<b>6.4</b>	9.0	<b>14.8</b>	10.8	<b>5.8</b>	<b>12.4</b>	<b>15.1</b>	<b>5.4</b>	<b>14.6</b>	8.4	<b>13.4</b>
2–4 long	<b>8.4</b>	<b>1.7</b>	<b>4.9</b>	5.4	<b>8.7</b>	6.5	<b>4.3</b>	<b>7.1</b>	<b>7.9</b>	<b>4.0</b>	<b>7.4</b>	5.9	<b>8.0</b>
5+ short	<b>18.3</b>	<b>5.0</b>	<b>16.0</b>	9.5	<b>24.8</b>	12.3	<b>10.2</b>	<b>17.2</b>	<b>20.8</b>	<b>27.0</b>	<b>22.3</b>	8.1	<b>13.4</b>
5+ long	<b>9.2</b>	<b>3.6</b>	<b>6.9</b>	6.2	<b>13.8</b>	7.9	<b>6.2</b>	<b>7.5</b>	<b>11.3</b>	<b>5.8</b>	<b>9.2</b>	4.9	<b>7.0</b>
Mother's age at birth													
20+	10.8	2.3	6.3	6.7	11.5	7.4	5.9	8.8	10.7	5.7	10.0	6.0	8.4
<20	11.3	2.7	5.7	5.9	9.3	9.4	5.3	8.3	11.8	3.2	10.7	6.5	9.6
Type of residence													
Urban	<b>8.7</b>	2.7	5.8	6.0	8.5	7.7	4.8	9.2	<b>8.0</b>	5.3	11.9	7.8	9.9
Rural	<b>12.5</b>	2.4	6.3	6.4	10.8	8.5	5.7	8.4	<b>12.5</b>	4.0	9.7	5.8	8.5
Mother's education													
Primary or higher	<b>8.5</b>	2.4	5.0	5.4	9.5	8.0	5.1	7.8	<b>9.0</b>	3.1	9.5	6.1	8.4
No	<b>13.0</b>	2.6	7.2	7.4	12.1	8.3	5.8	8.8	<b>13.0</b>	5.5	10.4	6.4	9.3
Father's education													
Primary or higher	10.2	2.3	5.1	<b>5.6</b>	9.4	7.4	4.9	8.4	<b>9.7</b>	4.0	9.1	6.3	8.4
No	12.5	3.0	<b>8.4</b>	<b>8.6</b>	10.7	9.4	5.8	8.8	<b>13.2</b>	5.3	10.6	5.9	9.9
Mother's occupation													
Other	8.9	2.7	5.5	7.1	11.7	8.6	4.6	8.9	11.0	<b>3.5</b>	9.8	6.4	9.4
Household	9.5	2.5	8.0	5.2	9.5	8.5	5.3	8.7	11.2	<b>4.2</b>	10.6	7.3	8.5
Agriculture	12.5	1.5	6.5	7.1	11.8	7.7	5.8	8.3	11.4	<b>13.0</b>	10.4	5.8	9.1
Father's occupation													
Other	<b>8.8</b>	2.2	5.1	5.6	9.8	7.7	4.9	8.5	<b>10.3</b>	3.8	10.0	7.4	8.9
Household	<b>10.9</b>	4.2	11.4	7.0	10.6	9.5	3.4	10.4	<b>11.0</b>	6.1	10.7	4.6	9.6

Continued

**Table 1** Continued

	DRCongo 2007	Egypt 2008	Ghana 2008	Kenya 2008/2009	Lesotho 2009/2010	Liberia 2007	Madagascar 2008/2009	Malawi 2010	Nigeria 2008	Sao Tome & Principe 2008/2009	Sierra Leone 2008	Tanzania 2009/2010	Zambia 2007
Agriculture	<b>12.8</b>	2.9	7.6	8.1	11.9	8.6	5.9	8.4	<b>12.4</b>	6.7	10.4	6.0	9.0
Region													
1	6.7	<b>3.4</b>	<b>3.9</b>	<b>6.5</b>	8.2	6.5	<b>5.2</b>	7.8	<b>10.3</b>	5.6	8.8	<b>4.4</b>	7.7
2	11.1	<b>1.7</b>	<b>8.4</b>	<b>6.3</b>	12.6	9.1	<b>7.5</b>	8.3	<b>12.8</b>	2.8	10.3	<b>4.7</b>	7.7
3	9.4	<b>2.9</b>	<b>4.8</b>	<b>5.9</b>	8.2	10.5	<b>4.1</b>	9.0	<b>13.5</b>	3.6	10.9	<b>2.9</b>	10.3
4	12.7	<b>2.4</b>	<b>3.0</b>	<b>5.3</b>	10.3	8.5	<b>5.8</b>		<b>11.1</b>	2.6	11.6	<b>6.3</b>	8.0
5	10.7		<b>5.8</b>	<b>10.6</b>	10.3	8.2	<b>5.0</b>		<b>9.8</b>			<b>7.7</b>	10.3
6	7.9		<b>6.5</b>	<b>3.7</b>	11.0	8.0	<b>4.7</b>		<b>6.6</b>			<b>4.7</b>	10.6
7	14.3		<b>4.3</b>	<b>7.0</b>	11.1							<b>8.9</b>	7.0
8	12.3		<b>9.4</b>	<b>6.5</b>	9.2							<b>8.1</b>	7.8
9	15.1		<b>3.5</b>		9.5							<b>5.2</b>	8.4
10	12.0		<b>11.8</b>		9.9							<b>3.5</b>	
11	10.2											<b>8.6</b>	
12												<b>6.2</b>	
13												<b>3.3</b>	
14												<b>5.8</b>	
15												<b>6.3</b>	
16												<b>7.8</b>	
17												<b>5.4</b>	
18												<b>6.6</b>	
19												<b>5.7</b>	
20												<b>12.3</b>	
21												<b>2.9</b>	
22												<b>5.2</b>	
23												<b>6.9</b>	


Reference categories used in the multivariate regression model are italicised. Proportions differing significantly ( $p < 0.01$ ,  $\chi^2$  test) are in bold.  
 Occupation other: Prof., Tech., Manag., Clerical, Sales, Services, Skilled manual, other. Household: Did not work, Household & Domestic, Unskilled manual, None. Agriculture: Agric-self-employed, Agric-employee.



**Figure 2** Wealth-related inequality in under-5 mortality versus average under-5 mortality in the 13 countries, Demographic and Health Surveys 2007–2010. \*Erreygers' corrected concentration index.<sup>19</sup>

overall inequality in under-5 mortality, except in Lesotho, Madagascar, Sao Tome & Principe and Sierra Leone where it was still substantial (>5%). Results obtained by Shorrocks decomposition of the variance were similar.

Further, the decomposition of wealth-related inequalities in under-5 mortality was applied to countries where C was significant. Here, unlike in figure 1, C had positive values because it measures inequality in predicted (ln odds) under-5 mortality, which has a negative mean. But in Sao Tome & Principe, as some children had a positive predicted value (a very high death probability), G and C measures were corrected (the minimum value was subtracted from all observations) and their sign is therefore inverted. The major determinants of wealth-related inequality in under-5 mortality differed in each country. In



**Figure 3** Trends in wealth-related inequality in under-5 mortality in DR Congo, Egypt, Madagascar, Nigeria and Sao Tome & Principe, Demographic and Health Surveys 2007–2010.

Egypt, inequality was mostly explained by household wealth itself, birth order and interval, father's education and father's occupation. In Madagascar, birth order and interval contributed mostly, followed by household wealth and skilled birth attendance. In Nigeria, inequality was mostly explained by the type and region of residence. In DR Congo and Sao Tome & Principe, apart from household wealth, mother's education and father's occupation were important determinants of wealth-related inequality. Determinants with a negative contribution (eg, rural residence, mother's low education and some regions) were more prevalent among poor households but negatively associated with under-5 mortality, thereby decreasing the wealth-related inequality.

Finally, determinants were ranked according to these contributions and rankings were summarised by determinants in figure 4. Considering the five countries together, determinants contributing most to overall inequality in under-5 mortality were: birth order and interval, region, mother's education and father's occupation. Determinants contributing most to wealth-related inequality in under-5 mortality were: household wealth, father's occupation and mother's education.

### DISCUSSION

In the context of increasing awareness about the role of socio-economic determinants in shaping health and health inequalities, the study investigated the relative contribution of determinants to overall inequality in under-5 mortality and to wealth-related inequality in under-5 mortality in a selection of African countries. In almost all countries, birth order and interval and region contributed the most to the overall inequality. In the five countries where wealth-related inequality was significant, household wealth, father's occupation and mother's education were overall important contributors, though differences were observed between countries.

The measure of C allowed the distinction of countries that require a whole population improvement of child health: Tanzania, Ghana, Kenya, Liberia, Malawi, Zambia, Sierra Leone and Lesotho and those that would also benefit from the addition of more poor-targeted interventions: Egypt, Madagascar, Sao Tome & Principe, DRCongo and Nigeria. C of under-5 mortality was reported for preceding DHS—Ghana 2003, Kenya 2003, Madagascar 1997, Malawi 2000, Nigeria 2003, Tanzania 2004 and Zambia 2001—by the World Bank.<sup>25</sup> In this report, Madagascar, Zambia, Kenya, Malawi and Ghana had a significant negative C. Though wealth-related inequality in under-5 mortality may have decrease since the last DHS, the difference in significance with results reported here can result from sample size: under-5 mortality was assessed until 10 years before the survey, whereas in the study at hand it was restricted to a 5-year period. A study using 1990–1999 DHS data from 60 countries showed a negative association between the level and inequality in under-5 mortality; countries with the highest level of under-5 mortality had a lower inequality.<sup>26</sup> In the study at hand, no significant relation between C\* and average under-5 mortality was observed, possibly explained by only using a limited set of African countries with most of these not showing under-5 mortality inequalities.

The regression analysis identified already known determinants of under-5 mortality in sub-Saharan Africa: household wealth, region, type of residence (rural/urban), parent's education, birth order and preceding birth interval and skilled birth attendance.<sup>8–15</sup> In addition, the analysis revealed an independent effect of the sex of the child; mortality was higher in males in Lesotho, Malawi, Nigeria and Sao Tome & Principe. This effect,

**Table 2** Multivariate logistic regression of under-5 mortality: regression coefficients and significance in the 13 countries, Demographic and Health Surveys 2007–2010

	DR Congo 2007	Egypt 2008	Ghana 2008	Kenya 2008/ 2009	Lesotho 2009/ 2010	Liberia 2007	Madagascar 2008/2009	Malawi 2010	Nigeria 2008	Sao Tome & Principe 2008/ 2009	Sierra Leone 2008	Tanzania 2009/ 2010	Zambia 2007
N (10 <sup>3</sup> )	8.3	10.7	2.7	5.4	3.5	4.6	11.3	19.4	26.9	1.8	4.9	7.6	5.6
Low household wealth	0.07	0.38	-0.58	0.02	0.02	0.09	0.07	-0.03	0.10	0.51	0.18	0.18	-0.20
No skilled birth attendance	0.20	0.01	-0.23	-0.13	0.34	0.02	0.13	0.07	-0.03	-0.57	0.31	0.06	0.00
Male	<b>0.24</b>	0.30	0.08	0.32	0.36	0.20	0.05	<b>0.29</b>	<b>0.13</b>	<b>1.28</b>	-0.17	0.29	0.20
Birth 2–4 short	0.00	<b>0.61</b>	-0.50	0.22	0.09	0.23	0.13	0.12	<b>0.44</b>	1.66	0.09	0.24	0.15
Birth 2–4 long	<b>-0.51</b>	-0.24	-0.44	-0.27	<b>-0.53</b>	-0.26	-0.16	<b>-0.50</b>	<b>-0.28</b>	1.07	<b>-0.56</b>	-0.25	<b>-0.46</b>
Birth 5+ short	0.42	0.80	0.78	0.01	0.49	0.41	<b>0.80</b>	0.41	<b>0.82</b>	<b>2.96</b>	<b>0.71</b>	0.08	0.00
Birth 5+ long	-0.40	0.47	-0.30	-0.36	-0.35	-0.03	0.22	<b>-0.54</b>	0.13	0.87	-0.25	-0.40	<b>-0.67</b>
Mother's age at birth <20	0.03	0.15	0.08	-0.30	-0.44	0.27	0.12	-0.20	<b>0.25</b>	-0.09	0.19	-0.10	-0.18
Rural residence	0.09	-0.07	0.19	0.00	0.13	-0.02	0.05	-0.12	<b>0.33</b>	-1.28	-0.51	-0.28	-0.35
Mother low education	0.34	-0.35	0.08	0.18	0.30	-0.25	-0.13	0.17	0.10	<b>1.56</b>	<b>-0.10</b>	0.00	0.12
Father low education	<b>-0.04</b>	0.17	0.36	<b>0.61</b>	-0.10	0.32	0.02	0.02	0.07	<b>-0.19</b>	0.22	0.01	0.23
Mother household	-0.02	-0.35	0.34	-0.36	-0.35	-0.19	0.17	-0.07	-0.14	0.05	-0.04	0.27	-0.11
Mother agriculture	0.14	-0.84	0.14	-0.19	-0.24	-0.49	0.03	-0.10	-0.07	<b>1.40</b>	0.12	-0.27	0.12
Father household	0.24	0.58	0.93	0.19	0.11	0.31	-0.41	0.23	0.07	1.20	0.09	-0.51	0.06
Father agriculture	0.22	0.28	0.44	0.37	0.12	0.11	0.06	0.01	-0.02	0.77	0.08	0.12	0.05
Region 2	0.18	<b>-0.76</b>	0.38	-0.11	<b>0.67</b>	0.27	0.29	0.05	0.17	-0.73	0.06	-0.27	-0.24
Region 3	-0.08	-0.40	0.25	-0.31	0.11	0.46	-0.28	0.16	0.18	-1.10	0.28	-0.63	0.36
Region 4	0.17	-0.49	-0.38	-0.50	0.27	0.18	0.04	0.19	0.19	-0.65	0.34	0.16	0.03
Region 5	-0.17		0.54	0.32	0.36	0.41	-0.30		0.01			0.51	0.18
Region 6	-0.33		0.35	-1.02	0.29	0.40	-0.19		-0.29			0.11	0.28
Region 7	0.41		-0.04	-0.24	0.34							0.65	-0.20
Region 8	0.22		0.61	-0.85	0.22							0.78	0.02
Region 9	0.43		-0.17		0.14							0.34	0.00
Region 10	0.07		0.73		0.08							-0.39	-1.97
Region 11	0.02											0.85	
Region 12												0.30	
Region 13												-0.27	
Region 14												0.29	
Region 15												0.55	
Region 16												0.54	
Region 17												0.34	
Region 18												0.58	
Region 19												0.31	
Region 20												<b>1.30</b>	
Region 21												-1.07	
Region 22												0.09	
Region 23												0.29	

Occupation other: Prof., Tech., Manag., Clerical, Sales, Services, Skilled manual, other. Household: Did not work, Household & Domestic, Unskilled manual, None. Agriculture: Agric-self-employed, Agric-employee.  
Significant regression coefficients (p<0.01) are in bold. To avoid overloading the table, CIs and exact p values are presented in an online supplementary appendix.

already observed at the univariate level in the respective DHS reports,<sup>7</sup> was also shown in a study assessing individual and community level effects on child mortality in 28 sub-Saharan African countries.<sup>27</sup> An excess male infant mortality was previously observed in high-income countries.<sup>28</sup> This male disadvantage was most likely explained by biological factors such as a lower resistance to infection, a higher risk of premature birth and a larger average body size and head circumference leading to difficult labour. A community-based

cohort study performed in rural Malawi documented a twofold mortality risk among 1-year-old to 2-year-old boys compared with girls of the same age.<sup>29</sup> In this subpopulation, possible explanations tended more towards biological (eg, malnutrition or sex-specific differences in vaccine effects) than behavioural (eg, preferential care) assumptions.

The analysis of the contribution of determinants to overall inequality in under-5 mortality allowed the determinants to be

**Table 3** Relative contribution (%) of determinants to overall inequality (G) and to wealth-related inequality (C) in under-5 mortality in the 13 countries, Demographic and Health Surveys 2007–2010

	G	Household wealth	Skilled birth attendance	Sex of the child	Birth order and interval	Mother's age at birth	Type of residence	Mother's education	Father's education	Mother's occupation	Father's occupation	Region
DR Congo	-0.128	2.4	7.3	5.3	35.6	0.4	3.4	17.4	-0.7	5.8	9.1	14.0
Egypt	-0.080	11.7	0.1	7.3	35.0	1.7	0.9	-2.4	2.8	3.0	12.2	27.9
Ghana	-0.108	1.9	0.2	0.6	22.1	0.5	2.2	1.9	12.5	5.6	16.3	36.3
Kenya	-0.129	0.1	0.2	5.8	7.2	1.9	0.0	3.0	15.8	5.7	7.5	52.6
Lesotho	-0.117	0.4	14.9	16.6	18.3	10.6	2.3	10.7	-0.5	7.0	1.9	17.6
Liberia	-0.082	1.4	0.2	7.4	30.2	11.2	-0.4	1.8	18.7	10.4	3.9	15.1
Madagascar	-0.068	2.5	6.0	0.5	47.6	-1.3	0.9	-1.3	0.5	-0.1	5.0	39.7
Malawi	-0.077	0.1	0.9	21.1	63.7	0.3	1.7	3.1	0.1	1.4	3.9	3.7
Nigeria	-0.124	5.0	-1.3	1.7	46.9	4.2	16.2	5.0	2.9	-0.5	-0.6	20.5
Sao Tome & Principe*	0.191	3.9	0.5	17.9	14.8	0.5	14.0	27.3	-1.0	6.6	8.9	6.6
Sierra Leone	-0.109	3.7	8.2	3.8	57.0	4.7	8.9	0.1	3.5	1.4	0.5	8.2
Tanzania	-0.102	-0.6	0.1	7.8	14.1	-1.0	6.7	0.0	0.0	8.9	4.3	59.8
Zambia	-0.092	4.6	0.0	6.7	46.4	-2.7	9.9	2.0	6.5	2.5	-0.1	24.1

	C	Household wealth	Skilled birth attendance	Sex of the child	Birth order and interval	Mother's age at birth	Type of residence	Mother's education	Father's education	Mother's occupation	Father's occupation	Region
DR Congo	0.057	13.1	11.3	-0.6	-1.2	0.4	12.5	33.1	-2.1	17.6	20.8	-4.9
Egypt	0.020	119.0	1.5	0.8	33.4	5.3	-13.8	-65.1	22.0	-25.3	49.5	-27.3
Madagascar	0.020	29.4	27.7	0.4	39.0	2.3	6.7	-25.4	5.0	-7.6	18.3	4.3
Nigeria	0.067	16.5	-3.2	0.3	11.1	4.9	33.5	11.6	6.9	-4.7	-2.3	25.4
Sao Tome & Principe*	-0.045	60.9	-12.3	2.0	4.5	-1.2	-43.4	76.3	-9.4	14.5	40.2	-32.2

The relative contributions were calculated by decomposing G and C using Wagstaff's method.<sup>20</sup>

\*As some children had a positive predicted value (a very high death probability) in Sao Tome & Principe, G and C measures were corrected (the minimum value was subtracted from all observations) and their sign is therefore inverted.  
C, concentration index; G, Gini index.


classified by importance. The relative importance methodology is increasingly used as a supplement to multiple regression analyses to better understand the role played by each predictor in a regression equation.<sup>30</sup> The analysis was performed by decomposing the Gini index. The same analysis was also performed by decomposing the variance using Shorrocks's method,<sup>23</sup> which gave similar results. Results could guide interventions aiming at reducing overall inequality in under-5 mortality by prioritising its determinants at the national level. For example, high parity and short-birth interval could be targeted through family planning in countries where birth order and interval explains most of the overall inequality in under-5 mortality. Increasing health promotion coverage and access to healthcare, for example, through decentralisation of health services, could be envisaged as a priority in countries where regional differences are the most important.

This is, to our knowledge, the first study investigating socioeconomic determinants of wealth-related inequality in under-5 mortality in Africa. In countries where wealth-related inequality was large, identifying determinants contributing the most to this inequality may help in prioritising interventions aiming at reducing socioeconomic inequality in under-5 mortality. Reducing wealth inequalities, for example, through community loans, would be relevant in countries where household wealth has a major and direct effect on inequality. But in countries like Sao

Tome & Principe or DR Congo, where mother's education explained most of the wealth-related inequality, the priority could be education and health promotion. The substantial contribution of the type and region of residence in Nigeria, independent of the other factors, may reflect inequality in healthcare access and could be more investigated. The role of father's occupation in wealth-related inequality could also be more investigated in countries where this determinant was important.

The study has a number of limitations. The transversal design of the survey does not allow the order of events to be taken into account (eg, the household was not necessarily in the same wealth quintile or region at the time of the event). Information of child death was collected retrospectively by interviewing mothers and therefore relies on the mother's will and ability to communicate the information. Other determinants reported elsewhere, for example, birth weight, breastfeeding, antenatal care, ethnicity or religion<sup>10 31 32</sup> were not included in models for analytical reasons (multicollinearity or high number of missing values). Age of children at the time of the survey was not taken into account; however, regression coefficients remained unchanged when adding theoretical age in the models. Finally, the interpretation of global patterns merging countries as diverse as Egypt and Nigeria in terms of national income and under-5 mortality must be done being mindful that the socioeconomic and political context plays a proximate role





**Figure 4** Ranking of contributions to wealth-related inequality versus overall inequality in under-5 mortality, five countries, 2007–2010. SBA, skilled birth attendance. Units are determinants' median rank (1=the first) across the five countries with a significant C: Egypt, Madagascar, Nigeria, DR Congo and Sao Tome & Principe.

in the process to inequalities as illustrated in the conceptual framework proposed by CSDH.<sup>6</sup>

## CONCLUSION

Generally, the main determinants of overall inequality in under-5 mortality were birth order and interval and region. The main determinants of wealth-related inequality in under-5 mortality were household wealth, father's occupation and mother's education. At the national level, the identified determinants contributing the most to either overall inequality or wealth-related inequality in under-5 mortality could serve as targets to improve both average child survival and equity.

### What is already known on this subject?

- ▶ Despite substantial progress in reducing average under-5 mortality, disparity in under-5 mortality remains high.
- ▶ Understanding the underlying determinants is essential to develop appropriate policy responses.

### What this study adds?

- ▶ The relative contribution of several determinants to overall inequality in under-5 mortality is given for 13 African countries.
- ▶ For five African countries with a significant wealth-related inequality in under-5 mortality, the relative contribution of the same determinants to this wealth-related inequality is given as well.
- ▶ Interventions aiming at reducing either overall under-5 mortality or socioeconomic inequality in under-5 mortality can therefore be prioritised.

**Acknowledgements** We thank the Measure Demographic and Health Survey programme for the datasets. We are grateful to Irene Ogali for her contribution to the application of Shorrocks' method and to Corey Sparks for his assistance with the creation of maps. We would like to express our thanks to the reviewers for their constructive and interesting comments.

**Contributors** NS designed the study. CVM analysed the data and wrote the manuscript. NS and HVO supervised the study and reviewed the manuscript.

**Collaborators** Irene Ogali contributed to the application of Shorrocks' method and Corey Sparks assisted with the creation of maps.

**Funding** This study is part of a PhD thesis financially supported by the Universit  Catholique De Louvain, Brussels, Belgium.

**Competing interests** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- 1 UN Inter-agency Group for Child Mortality Estimation. *Levels & trends in child mortality*. United Nations Children's Fund; 2011, Report 2011.
- 2 Commission on Social Determinants of Health WHO. *Closing the gap in a generation. Health equity through action on the social determinants of health*. Geneva, Switzerland: Commission on Social Determinants of Health; 2008, final report.
- 3 Houweling TA, Kunst AE. Socio-economic inequalities in childhood mortality in low- and middle-income countries: a review of the international evidence. *Br Med Bull* 2010;93:7–26.
- 4 Speybroeck N, Konings P, Lynch J, et al. Decomposing socioeconomic health inequalities. *Int J Public Health* 2010;55:347–51.
- 5 Hosseinpoor AR, van Doorslaer E, Speybroeck N, et al. Decomposing socioeconomic inequality in infant mortality in Iran. *Int J Epidemiol* 2006;35:1211–19.
- 6 World Health Organisation. A conceptual framework for action on the social determinants of health. Social Determinants of Health, 2010, Discussion paper 2.
- 7 Measure DHS. *The demographic and health surveys program*. Macro International, 2013. <http://www.measuredhs.com> (accessed 26 Feb 2013).
- 8 Houweling TA, Caspar AE, Looman WN, et al. Determinants of under-5 mortality among the poor and the rich: a cross-national analysis of 43 developing countries. *Int J Epidemiol* 2005;34:1257–65.
- 9 Houweling TA, Kunst AE, Moser K, et al. Rising under-5 mortality in Africa: who bears the brunt? *Trop Med Int Health* 2006;11:1218–27.
- 10 Antai D. Inequalities in under-5 mortality in Nigeria: do ethnicity and socioeconomic position matter? *J Epidemiol* 2011;21:13–20.
- 11 Antai D. Regional inequalities in under-5 mortality in Nigeria: a population-based analysis of individual- and community-level determinants. *Popul Health Metr* 2011;9:6.
- 12 Antai D, Wedren S, Bellocco R, et al. Migration and child health inequities in Nigeria: a multilevel analysis of contextual- and individual-level factors. *Trop Med Int Health* 2010;15:1464–74.
- 13 Tottrup C, Tersbol BP, Lindeboom W, et al. Putting child mortality on a map: towards an understanding of inequity in health. *Trop Med Int Health* 2009;14:653–62.
- 14 Anyamele OD. Urban and rural differences across countries in child mortality in sub-Saharan Africa. *J Health Care Poor Underserved* 2009;20:90–8.
- 15 Lykens K, Singh KP, Ndukwe E, et al. Social, economic, and political factors in progress towards improving child survival in developing nations. *J Health Care Poor Underserved* 2009;20:137–48.
- 16 Ayotunde T, Mary O, Melvin AO, et al. Maternal age at birth and under-5 mortality in Nigeria. *East Afr J Public Health* 2009;6:11–14.
- 17 Adetunji JA, Shelton JD. Ethical issues in the collection, analysis and dissemination of DHS data in sub-Saharan Africa. 2011. <http://uaps2011.princeton.edu/papers/110641> (accessed 26 Feb 2013).
- 18 Kakwani N, Wagstaff A, van Doorslaer E. Socioeconomic inequalities in health: measurement, computation, and statistical inference. *J Econometrics* 1997;77:87–103.
- 19 Erreygers G. Correcting the Concentration Index. *J Health Econ* 2009;28:504–15.
- 20 Wagstaff A, van Doorslaer E, Watanabe N. On decomposing the causes of health sector inequalities with an application to malnutrition inequalities in Vietnam. *J Econometrics* 2003;112:207–23.
- 21 Wagstaff A, van Doorslaer E. Overall versus socioeconomic health inequality: a measurement framework and two empirical illustrations. *Health Econ* 2004;13:297–301.
- 22 Van de Poel E, Hosseinpoor AR, Jehu-Appiah C, et al. Malnutrition and the disproportional burden on the poor: the case of Ghana. *Int J Equity Health* 2007;6:21.
- 23 Shorrocks AF. *Decomposition procedures for distributional analysis: a unified framework based on the Shapley Value*. University of Essex and Institute for Fiscal Studies, 1999.
- 24 World Health Organisation. *Health systems financing: the path to universal coverage*. Geneva: WHO; 2010, The World Health report.

- 25 Gwatkin DR, Rutstein S, Johnson K, *et al.* Socio-economic differences in health, nutrition, and population within developing countries. Country reports on HNP and poverty. 2007.
- 26 Wang L. *Determinants of child mortality in low-income countries: empirical findings from demographic and health surveys*. The World Bank, 2002.
- 27 Boco AG. Individual and community level effects on child mortality: an analysis of 28 demographic and health surveys in sub-Saharan Africa. DHS working papers MEASURE DHS. 2010.
- 28 Drevenstedt GL, Crimmins EM, Vasunilashorn S, *et al.* The rise and fall of excess male infant mortality. *Proc Natl Acad Sci USA* 2008;105:5016–21.
- 29 Ashorn P, Maleta K, Espo M, *et al.* Male biased mortality among 1–2 year old children in rural Malawi. *Arch Dis Child* 2002;87:386–7.
- 30 Tonidandel S, LeBreton JM. Relative importance analysis: a useful supplement to regression analysis. *J Bus Psychol* 2011;26:9.
- 31 Antai D, Antai J. Individual- and contextual-level determinants of social inequities in under-five mortality in Nigeria: differentials by religious affiliation of the mother. *World Health Popul* 2009;10:22–36.
- 32 Kayode GA, Adekanmbi VT, Uthman OA. Risk factors and a predictive model for under-five mortality in Nigeria: evidence from Nigeria demographic and health survey. *BMC Pregnancy Childbirth* 2012;12:10.