

THEORY AND METHODS

Conceptualisation, development, and evaluation of a measure of unplanned pregnancy

G Barrett, S C Smith, K Wellings

J Epidemiol Community Health 2004;58:426–433. doi: 10.1136/jech.2003.014787

Study objective: To develop a measure of unplanned pregnancy that is valid, reliable, and appropriate in the context of contemporary demographic trends and social mores and can be used in a variety of situations, including the production of population prevalence estimates.

Design: A two stage study design: qualitative (inductive) methods to delineate the construct of pregnancy planning, and quantitative/psychometric methods to establish the means of measurement.

Setting: Eight health service providers (comprising 14 clinics, including antenatal, abortion, and one general practitioner) across London, Edinburgh, Hertfordshire, Salisbury, and Southampton in the UK.

Participants: Samples comprised a mixture of pregnant (continuing pregnancy and opting for abortion) and recently pregnant (post-abortion and postnatal) women. At the qualitative stage, 47 women took part in depth interviews (20 of whom were re-interviewed after the birth of their baby). Items were pre-tested with 26 women, and two psychometric field tests were carried out with, respectively, 390 and 651 women.

Main results: A six item measure of unplanned pregnancy was produced. Psychometric testing demonstrated the measure's high reliability (Cronbach's $\alpha=0.92$; test-retest reliability=0.97) and high face, content, and construct validity. Women's positions in relation to pregnancy planning are represented by the range of scores (0–12).

Conclusions: A psychometric measure of unplanned pregnancy, the development of which was informed by lay views, is now available. The measure is suitable for use with any pregnancy regardless of outcome (that is, birth, abortion, miscarriage) and is highly acceptable to women.

See end of article for authors' affiliations

Correspondence to:
Dr G Barrett, Department of Health and Social Care, Brunel University, Borough Road, Middlesex TW7 5DU, UK; geraldine.barrett@brunel.ac.uk

Accepted for publication 10 December 2003

The concept of "unplanned pregnancy" is widely used in health research and policy. Attempts to measure it have been numerous, varying from studies in which the concept is assumed to be self evident to those in which more sophisticated measurement strategies have been used.^{1–5} The approach taken by large national surveys has tended towards the latter, eliciting planning status by means of multi-dimensional questions, probing (in various combinations) intentions, contraceptive use, reactions to pregnancy, timing of pregnancy plans, and family size intentions.^{6–12} The most influential of these surveys has been the US National Survey of Family Growth¹² whose forms of measurement have been widely adopted.^{13–15}

In recent years, however, there has been growing awareness of the limitations of existing questions.^{5 16–21} No new estimates of unplanned pregnancy have been produced in Britain since 1991, and in the US National Survey of Family Growth items have been added incrementally to ensure validity.^{12 22} As most questions currently used to assess unplanned pregnancy were developed several decades ago, before legal abortion was available and when the primary concern was with excess fertility in marital relationships,²⁰ it is not surprising that such questions are becoming dated. Also, as therapeutic advances have effectively raised the upper age limit for pregnancy and as employment opportunities have increased for women, the social context of childbearing has changed. Furthermore, measurement to date has tended to assume congruence between intentions and behaviour despite evidence to the contrary.^{4 6 7 17}

Calls are now being made for a reconsideration of the conceptual basis of unplanned pregnancy.^{17–19} Improvements to the yield of national fertility surveys will only come, it is said, from "intensive work to refine the measurement of often elusive concepts".¹⁰ In this spirit we began a three year

study in 1998 to develop a new measure of unplanned pregnancy for use in Britain. We describe here the development and psychometric validation of the measure.

METHODS

The overall aim of the study was to develop a measure of pregnancy planning/intention that is valid, reliable, and appropriate in the context of contemporary demographic trends and social mores, and can be used to establish population estimates of unplanned pregnancy. To achieve this we used a two stage study design: (1) qualitative methods to delineate the construct of pregnancy planning; and (2) quantitative/psychometric methods to establish the means of measurement.

Conceptualisation

To develop a conceptual model of pregnancy planning/intention, we used depth interviews (that is, flexible interviews that use normal modes of conversation) to elicit women's accounts of the circumstances in which they became pregnant. The aim was to build a model based on the key elements of the interviews through which an understanding of women's experiences could be gained. (Details of the methodology and qualitative findings have previously been reported.^{23 24})

As previous evidence had suggested that answers elicited before and after birth may be different,^{25 26} we conducted follow up interviews with women who continued their pregnancies after they had their babies. The aim was to assess if, or how, women's accounts changed over this time period

Item development and piloting

The conceptual model produced during the qualitative stage informed item development; items were developed, without

limit, until the dimensions of the model were adequately represented. The items were piloted with a small sample of women and qualitative interviews were used to check women's understanding. Amendments to the items were made incrementally during piloting.

Item analysis and selection (first psychometric field test)

We screened items for homogeneity²⁷ using inter-item correlations and Cronbach's α .²⁸ We then devised a five step strategy for item analysis and selection: (1) Remove items with more than 5% missing data²⁹; (2) Remove any item with a maximum endorsement frequency of $\geq 80\%$ on any response option^{30 31}; (3) Remove any item with an item-total correlation of < 0.2 ^{30 31}; (4) Rank the remaining items according to the item-total correlations and then, starting with the lowest rank, remove items if they correlate highly (that is, > 0.75) with another question^{27 30-32}; (5) Return items to the scale, in reverse order of removal, until an α of > 0.90 is reached.^{27 32} (Although a criterion of 0.7 is often cited for internal consistency, we used a more stringent criterion to allow for the possibility that the Cronbach's α might be lower in future samples.^{27 32}) The strategy enabled us to maximise homogeneity while still maintaining content validity.

Evaluating the item reduced measure (second psychometric field test)

A second independent field test was carried out to establish the psychometric properties of the final, item reduced, measure (appendix, available to view on the journal web site <http://www.jech.com/supplemental>).

Before analysis, missing data from this field test were imputed for the 18 women with missing data using the method applied to the SF-36³³ (that is, where a subject has completed at least 50% of items of a scale, the mean score of their completed items can be substituted for the missing items).

Acceptability of the measure

Acceptability was assessed by examining rates of missing data for the overall score and the distribution of scores. The reading level of the measure was assessed using the Flesch-Kincaid grade level scale.³⁴ Field notes were kept by researchers of women's experiences of completing the measure; notes were based on the researchers' observations and informal questioning of women in clinics.

Reliability

Internal consistency was assessed using the Cronbach's α statistic (> 0.7 indicating acceptable reliability^{28 35}), and test-retest reliability was examined in two ways: (1) a standard test-retest where a sub-sample of women were required to complete the repeat measure 7 to 14 days after first completion (an interval comparable to that used for other measures³¹); and (2) a long term test-retest that only included women who had completed the measure initially when they were pregnant, and then completed the repeat measure some months later, after the birth. The rationale for this second test was to assess the stability of scores before and after birth in light of existing evidence which suggests that women's reporting is unstable over this period.^{25 26} In both instances, test-retest reliability was measured using the weighted κ (the non-parametric equivalent of the intra-class correlation coefficient), a score of 0.61–0.80 indicating "substantial" reliability and > 0.80 indicating "almost perfect" reliability.³⁶

Validity

Content validity was assessed by comparing items in the final item reduced instrument with the conceptual model.

Two methods were used to assess construct validity: principal component analysis for within scale analyses; and hypothesis testing. We used principal component analysis (using varimax rotation and requesting as many factors as there were eigenvalues > 1) to test the hypothesis that all variables would load onto one factor.³⁷ For hypothesis testing there were two levels of hypotheses, from the qualitative findings and from the literature (table 1).

Although construct validity is also sometimes tested by considering the relation between the new measure and an established measure of a similar construct (convergent validity) or with a known measure of a different construct (discriminant), not enough is known about the nature of the construct of "pregnancy planning" to enable these hypotheses to be formulated. Criterion related validity is usually established by comparing a new psychometric measure with an established measure (ideally a "gold standard") of the same construct. However, as the absence of an existing measure was the reason for developing this measure, testing (concurrent or predictive) criterion related validity was also not possible. Responsiveness refers to an instrument's ability to detect change (over time) in a dynamic construct (for example, change in health status); as conception is an event at one point in time and therefore not a dynamic construct, testing responsiveness was not appropriate.

Interpreting the scores

Interpretation of scores is normally an ongoing process over the life of a measure. As a first step in this process, we used a content based method of interpretation,³⁸ using the item score patterns in the second field test and data from the qualitative stage, to provide the contextual detail necessary for initial interpretation of the scores.

Sampling

As we aimed to develop a measure that could be used to produce population estimates, women who were (or had been) pregnant were our target population. Although many men clearly have an important role in pregnancy planning, we did not include them on the grounds that not all men know about the pregnancies of their partner (or ex-partner), and because relying on information collected from couples, rather than women alone, would introduce substantial biases into a population sample.

All samples were constructed to include women of all ages whose pregnancies were continued and those which ended in abortion. Recruitment of study participants was from eight health service providers (comprising 14 clinics, including antenatal, abortion, and one general practitioner) across London, Edinburgh, Hertfordshire, Salisbury, and Southampton in the UK.

The qualitative sample was purposively sampled according to the above criteria, and for follow up, all women who were eligible and could be contacted were re-interviewed. The pilot/item development sample was also purposively sampled.

As psychometric field test samples must reflect the populations for whom the measure is designed,^{27 30} unselected clinic populations were invited to take part until the ratio of abortions to live births in the samples was consistent with that in the national population³⁹⁻⁴² (that is, abortions comprise 22% of conceptions). Women who were continuing their pregnancy or opting for abortion were recruited in hospital clinics. Three researchers (led by GB) carried out the recruitment. Women were approached directly by the researcher in a waiting room or a side room at some time during their appointment; only occasionally women declined

Table 1 Construct validity hypothesis tests

Hypothesis	Variable	Score range (median)	Mean rank	Significance test and p value	
Level 1: Strong hypotheses from qualitative findings:					
Higher scores will be associated with continued pregnancies and lower scores with pregnancies ending in abortion.	Pregnancy outcome: continued pregnancy	0–12 (11)	393.53	Mann-Whitney U, p<0.0001	
		0–12 (2)	98.48		
“Living with husband” status will be associated with higher scores, other categories associated with lower scores.	Marital status/live with: husband	0–12 (11)	431.83	Kruskal-Wallis, p<0.0001	
		0–12 (6)	283.25		
		0–12 (2)	145.68		
Older age will be associated with higher scores (although the full range of scores likely on all ages).	Age group: not husband or partner	<20	154.90	Kruskal-Wallis, p<0.0001	
		20–24	240.18		
		25–29	307.57		
		30–34	398.96		
		35–39	397.37		
		40+	365.29		
Higher educational status will be associated with higher scores (although full range of scores likely for all levels of educational status).	Educational level: school	0–12 (6)	278.92	Jonckheere-Terpstra, p<0.0001	
		0–12 (7)	297.97		
		0–12 (11)	367.32		
Level 2: Hypotheses from the literature, neither proved nor disproved by qualitative findings:					
Black women will have lower scores.	Ethnicity: White British	0–12 (10)	335.59	Kruskal-Wallis, p=0.013	
		White other	1–12 (10)		331.37
		Asian/Asian British	1–12 (10)		331.63
		Black/Black British	0–12 (5)		246.75
		Mixed/other	1–12 (8)		272.32
The second child (actual births) will have the highest scores, and the third-plus child lowest scores.	Child order: Continued pregnancies only:	was/will be first child	1–12 (10)	Kruskal-Wallis, p<0.0001	
		was/will be second child	1–12 (11)		
		was/will be third or more child	0–12 (9)		
			197.32		

to participate. Postnatal women were recruited in two ways. Firstly, women (14 in first field test and 170 in second field test) were identified from records of recent births at the participating hospitals and were sent the questionnaire via post; response rates were 79% (11) and 67% (112) respectively. Secondly, postnatal women were recruited at community clinics run by health visitors. Sample sizes complied with guidance.⁴³

For the test-retest samples, women were invited to volunteer to complete a second questionnaire at home. Because of issues of confidentiality (particularly the problem of sending material about pregnancy to women's homes), women undergoing abortion were excluded from this process, and therefore from the standard test-retest. Of the 467 women invited to participate, 340 (73%) agreed. For both test-retests, 121 women were selected (on a quota basis to include a range of ages) to achieve the sample sizes necessary for repeated observations.⁴⁴

Ethical approval

Multi-centre ethical approval was obtained for the study.

RESULTS

Samples

The qualitative sample comprised 47 women: 28 were continuing their pregnancies (although one had a miscarriage a couple of days before the interview), two were about to have abortions, and 17 had recently had abortions, most within the past two weeks. Women's ages ranged from 15 to 43. Women's educational and occupational levels and marital/relationship situations varied widely.

Of the 27 women in the qualitative sample who subsequently had a baby, 20 were re-interviewed. One declined an

interview and six could not be recontacted. The mix of ages and personal circumstances in the sample was, however, maintained. At the time of follow up, the infants' ages ranged from two to six months, and the time between interviews was seven to ten months.

Twenty six women, aged 16 to 42, took part in the piloting of the items. Seventeen were continuing their pregnancies, five were about to have abortions, and six had had babies in the past three months.

Altogether 390 women took part in the first field test and 651 in the second field test. Table 2 shows the characteristics of both field test samples. The samples were consistent with national data in terms of birth/abortion ratio and were largely consistent in terms of age distribution and marital status, although women born abroad were slightly over-represented^{45–47} (table 3).

Ninety eight women (81%) completed the repeat measure for the standard test-retest; 90 (74%) were in the seven to 14 day window eligible for analysis.

Ninety women (76%) completed the repeat measure for the long term test-retest; 87 (72%) were eligible for analysis. (Two women had become pregnant again and one woman was still pregnant at 39 weeks). The interval between the two questionnaires was six plus months for most women.

Conceptualisation

The circumstances in which women became pregnant are summarised by six thematic areas: (1) expressed intentions; (2) desire for motherhood; (3) contraceptive use; (4) pre-conceptual preparations; (5) personal circumstances/timing; and (6) partner influences. These areas formed the dimensions of the conceptual model (fig 1). The model reflects the complexities of women's accounts by

Table 2 Characteristics of the women in the psychometric field test samples

Variable	Field test 1		Field test 2	
	%	(n)	%	(n)
Pregnancy situation	n=390		n=651	
continuing pregnancy*	63.8	(249)	47.3	(308)
abortion†	18.5	(72)	22.9	(149)
postnatal‡	17.7	(69)	29.8	(194)
Age	n=385		n=648	
under 20	7.0	(27)	11.8	(77)
20–24	13.5	(52)	15.6	(101)
25–29	24.4	(94)	21.0	(136)
30–34	29.6	(114)	30.1	(195)
35–39	20.8	(80)	17.7	(115)
40+	4.7	(18)	3.7	(24)
range	14–47		14–47	
Number of children	n=388		n=651	
0	37.4	(145)	37.5	(244)
1	34.8	(135)	36.1	(235)
2	19.1	(74)	17.8	(116)
3	5.7	(22)	5.7	(37)
4	2.1	(8)	1.5	(10)
5+	0.8	(4)	1.4	(9)
Who women live with	n=387		n=651	
husband	55.6	(215)	50.4	(328)
partner	20.9	(81)	19.8	(129)
husband and parents	2.6	(10)	2.0	(13)
partner and parents	1.8	(8)	2.3	(15)
parents	5.1	(20)	9.5	(62)
alone	9.5	(37)	10.0	(65)
other relatives or friends	4.1	(16)	4.3	(28)
other	0.3	(1)	1.7	(11)
Woman's place of birth	n=388		n=648	
Britain	73.5	(285)	77.6	(503)
elsewhere	26.5	(103)	22.4	(145)
Ethnicity	n=389		n=650	
White				
white British	65.0	(253)	71.7	(466)
white Irish	3.1	(12)	2.3	(15)
white other	9.5	(37)	8.9	(58)
Mixed				
mixed—white and black Caribbean	0.5	(2)	0.6	(4)
mixed—white and black African	0.5	(2)	0.2	(1)
mixed—white and Asian	0.5	(2)	0.6	(4)
mixed-other	0.8	(3)	0.3	(2)
Asian				
Asian—Indian	2.1	(8)	1.4	(9)
Asian—Pakistani	1.8	(7)	4.5	(10)
Asian—Bangladeshi	1.3	(5)	0.8	(5)
Asian—other	2.6	(10)	1.7	(11)
Black/Black British				
Black Caribbean	4.4	(17)	3.5	(23)
Black African	5.4	(21)	4.0	(26)
Black other	0.3	(1)	0.3	(2)
Chinese or other				
Chinese	1.5	(6)	0.8	(5)
Other	0.8	(3)	1.4	(9)

*Field test 2 includes one woman who was continuing her pregnancy but miscarried in the 24 hours before completing the measure. †In field tests 1 and 2, 68 (94%) and 147 (99%) respectively were currently pregnant and about to terminate their pregnancies. ‡In field tests 1 and 2, 54 (78%) and 177 (91%) respectively had a child under 1 year.

encompassing a range of positions on each dimension (for example, positive, negative, ambivalent) and by neither requiring, nor assuming, congruence between dimensions.

In the follow up interviews, women were, overall, extremely consistent in their descriptions of the circumstances in which they became pregnant; confirming many features of their earlier interviews spontaneously. Only one woman modified an aspect of her account (her contraceptive use) relating to the conceptual model. (In contrast, greater change was noted regarding decision making after confirmation of pregnancy, and the interviews showed that women could clearly distinguish between their thoughts and feelings about events leading to their pregnancies and their thoughts and feelings about these events later, in light of their new experiences.)

Development and piloting of items

Eleven items were developed from the conceptual model: contraceptive use was represented by two items, personal circumstances/timing by four, partner influences by two, and the remaining dimensions by one item each. During piloting, one item (relating to contraception) was separated into two items, and minor changes to the wording and layout were made.

Item analysis and selection

No item failed the threshold of >5% missing data. One question failed the criterion of an endorsement frequency of over 80% on any one response option, hence was removed. Table 4 shows the inter-item and item-total correlations of the remaining items. No item had an item-total correlation of <0.2, and

Table 3 Comparison of field test samples with national data

	Field test 1 sample		Field test 2 sample		England and Wales, 2000*	
Age	Women opting for abortion (n=71):		Women opting for abortion (n=146):		Age at abortion (n=175542):	
	%	(n)	%	(n)	%	(n)
under 20	15.5	(11)	24.7	(36)	21.1	(36966)
20-24	29.6	(21)	26.7	(39)	26.8	(47099)
25-29	29.6	(21)	28.1	(41)	21.6	(37852)
30-34	15.5	(11)	9.6	(14)	16.4	(28735)
35-39	5.6	(4)	8.2	(12)	10.6	(18589)
40+	4.2	(3)	2.7	(4)	3.6	(6253)
Age	Women continuing pregnancy (n=247)		Women continuing pregnancy (n=307):		Age of mother at live birth (n=604441):	
	%	(n)	%	(n)	%	(n)
under 20						
20-24	5.7	(14)	11.4	(35)	7.6	(45846)
25-29	10.9	(27)	14.3	(44)	17.8	(107741)
30-34	24.7	(61)	18.6	(57)	28.2	(170701)
35-39	31.6	(78)	32.2	(99)	29.8	(180113)
40+	23.5	(58)	20.2	(62)	14.1	(84974)
	3.6	(9)	3.3	(10)	2.5	(15066)
	Women continuing pregnancy/postnatal women (n=311): proportion who were married:		Women continuing pregnancy/postnatal women (n=500): proportion who were married:		Proportion of live births registered by married parents:	
Total	%	(n)	%	(n)	%	(n)
	66.6	(207)	64.4	(322)	60.5	(365836)
By age group						
under 20	0	(0)	9.8	(4)	10.3	(4742)
20-24	32.3	(10)	29.5	(18)	37.4	(40262)
25-29	63.0	(46)	66.0	(62)	65.4	(111606)
30-34	78.0	(78)	76.2	(138)	75.6	(136165)
35-39	81.6	(62)	84.5	(87)	73.8	(62671)
40+	73.3	(11)	65.0	(13)	69.0	(10390)
	Women continuing pregnancy/postnatal women (n=316): proportion who were born in the UK:		Women continuing pregnancy/postnatal women (n=499): proportion who were born in the UK:		Proportion of births registered by women who were born in the UK:	
Total	%	(n)	%	(n)	%	(n)
	76.9	(243)	79.4	(396)	84.5	(510835)

*Office for National Statistics 2001^{45, 46}; figures for Scotland similar but not shown.

applying the strategy of considering the inter-item correlations resulted in eight items (2, 4a, 4b, 4c, 4d, 6, 7, 8) being removed. Cronbach's α of the remaining three items was 0.78. To achieve an α of >0.90, three items (4b, 6, 7) were returned.

Acceptability

After imputation of missing data, scores were available for all women in the sample. All scores were represented; 2.3% of women scored the minimum (0) and 25.0% the maximum




Figure 1 Conceptual model of pregnancy planning/unplanned pregnancy.

Key points

- A new, psychometrically evaluated, measure of unplanned pregnancy is now available for use.
- The new measure is based on lay views, rather than professional conceptualisations, of pregnancy planning.
- The measure is suitable for use with all women regardless of (intended or actual) pregnancy outcome.

(12). The skew statistic was -0.4 , however visual inspection of the distribution suggested that the scores were negatively skewed, possibly bimodal (with peaks at scores 2 and 12). The readability level of the measure was 6.7 on the Flesch-Kincaid grade level score (that is, suitable for an 11 year old), and most women completed the measure in 60–90 seconds. The measure was well received and did not cause offence.

Reliability

The Cronbach’s α was 0.92. (Item-total correlations ranged from 0.60 to 0.89 and inter-item correlations ranged from 0.44 to 0.83.) For the standard test-retest the weighted κ was 0.97, and for the long term test-retest it was 0.86.

Validity

Comparison of the six item measure with the conceptual model showed that content validity had been maintained, with one question representing each dimension of the model. All hypotheses to test construct validity were supported (table 1), suggesting that the scale is indeed measuring the degree to which a pregnancy is planned. The results of principal component analysis confirmed that all variables loaded onto one factor (eigenvalue 4.33), with high factor loadings for each item: qu1-0.70; qu2-0.90, qu3-0.93, qu4-0.90, qu5-0.89, qu6-0.75.

Interpreting the scores

The increasing scores of the measure (zero to 12) represent increasing degrees of pregnancy planning/intention and there are no obvious cut points in the scale; each score provides additional information. In terms of producing population estimates, we suggest (on the basis of preliminary interpretation) the division of scores into a minimum of three groups—that is, 10–12 (planned), 4–9 (ambivalent); and 0–3 (unplanned).

DISCUSSION

We developed a six item measure of unplanned pregnancy. Psychometric testing demonstrated the high internal consistency, high stability (standard and longer term), and excellent face, content, and construct validity of the measure. One limitation is that for reasons of confidentiality, the standard test-retest only included women who continued their pregnancies, thus our assessment of test-retest reliability does not provide any information about the stability of the scale when used with women whose pregnancies ended in abortion. Interestingly, the findings of the long term test-retest (and the qualitative follow up interviews) directly contradict previous evidence concerning the stability of women’s reports of pregnancy planning after birth.^{25 26} The reason for this may be that the items of the measure permit women a wider range of answers and therefore do not force women into categories that may be invalid. The measure was developed in Britain and is therefore appropriate for use with this population. As with other measures, re-validation would be required before application to other countries.

Compared with previous questions used to assess pregnancy planning, the measure has a number of advantages: it makes no assumptions about the nature of women’s relationships; it does not rely on women having fully formed childbearing plans; it does not assume a particular form of family building; and it is suitable for use with any pregnancy regardless of outcome. Because of its conceptual basis, the measure does not presume that women have clearly defined intentions and/or behaviour consistent with intentions. Women may occupy a range of positions in relation to pregnancy planning, and these are represented by the range of scores from zero to 12. The scores also provide more sophisticated information about pregnancy planning than the

Policy implications

- Unplanned pregnancy is often used as a proxy indicator of poor sexual health, and reducing the number of unplanned pregnancies is a policy aim of many countries around the world, including the USA and the UK.
- Existing methods of eliciting pregnancy planning status have become dated.
- A new measure will facilitate the production of reliable estimates of unplanned pregnancy.

Table 4 Inter-item and item-total correlations

Inter-item correlations:										
q1 (1)*	1.0									
q2	0.7924	1.0								
q4a	0.4792	0.4636	1.0							
q4b (2)	0.4998	0.4228	0.7811	1.0						
q4c	0.5268	0.4858	0.7503	0.7481	1.0					
q4d	0.4328	0.4164	0.7458	0.7064	0.8317	1.0				
q5 (3)	0.6080	0.5808	0.7172	0.7453	0.6815	0.6633	1.0			
q6 (4)	0.5170	0.4688	0.7317	0.8140	0.6870	0.6859	0.8137	1.0		
q7 (5)	0.5292	0.4955	0.7225	0.7592	0.7285	0.6670	0.8042	0.7645	1.0	
q8	0.4807	0.4395	0.7025	0.6934	0.6784	0.6303	0.6976	0.7259	0.7824	1.0
q9 (6)	0.4134	0.3790	0.5125	0.5351	0.4773	0.4646	0.6241	0.5464	0.5564	0.5446
	q1(1)	q2	q4a	q4b(2)	q4c	q4d	q5(3)	q6(4)	q7(5)	q8
Item-totals†	0.6369	0.5915	0.8210	0.8354	0.8157	0.7716	0.8641	0.8429	0.8498	0.7892
										q9(6)
										0.6165

*Figures in parentheses correspond to item numbers of the measure (appendix, available to view on the journal web site <http://www.jech.com/supplemental>).
†An item-total correlation is the correlation of the individual item with the scale total omitting that item.

dichotomous categories of planned and unplanned. The measure is short (only six items) and highly acceptable (that is, easy to understand, inoffensive, and quick to complete), attributes that make it suitable for use in large scale surveys.

The measure, with its conceptual basis, represents a clear break with the forms of measurement found in the previous British surveys and the current US and Demographic and Health Surveys (the last being the main data source of the international family planning movement). As such, the measure avoids the assumption that members of modern (post-demographic transition) societies are universally rational and instrumental in terms of their fertility decisions and control; an assumption that some have seen as characterising research on fertility and fertility change in the 20th century.^{48, 49} Instead, the measure permits representation of a range of positions (for example, actions congruent with intentions, actions inconsistent with intentions, ambivalence in fertility intentions and actions, etc), thereby providing a more complex and realistic of portrayal of human fertility behaviour than existing questions.

ACKNOWLEDGEMENTS

We wish to acknowledge the valuable contribution of our project collaborators: Lynne Cho, Karen Dunnell, Penny Edgington, Elaine Fisher, Anna Glasier, Isaac Manyonda, Stanley Okolo, Catherine Paterson, Connie Smith, and R William Stones. We would also like to thank the many others who have helped with this project, including: Maureen Batley, Audrey Brown, Jill Brown, Jenny Charman, Carrie Free, Rolla Khadduri, Patricia Kingori, Sunethra Kossine, Maya Malalgoda, Janet Peacock, Jan Sanders, Margaret Thorogood, Ros Tolcher, and Maddy Ward.



The appendix is available to view on the journal web site (<http://www.jech.com/supplemental>).

Authors' affiliations

G Barrett, K Wellings, Centre for Sexual and Reproductive Health Research, Department of Public Health and Policy, London School of Hygiene and Tropical Medicine, University of London, London, UK
S C Smith, Health Services Research Unit, Department of Public Health and Policy, London School of Hygiene and Tropical Medicine, University of London

Funding: this project was supported by a Medical Research Council/London Region special training fellowship in Health Services Research

Conflicts of interest: none declared.

REFERENCES

- Jurich J. The relationship of modernity of sex roles to pregnancy planning. *Sociological Focus* 1984;**17**:223–42.
- Najman JM, Morrison J, Williams G, et al. The mental health of women six months after they give birth to an unwanted baby: a longitudinal study. *Soc Sci Med* 1991;**32**:241–7.
- Miller WB. Reproductive decisions: how we make them and how they make us. *Advances in Population: psychological perspectives* 1994;**2**:1–27.
- Harris K, Campbell E. The plans in unplanned pregnancy: secondary gain and the partnership. *Br J Med Psychol* 1999;**72**:105–20.
- Morin P, Payette H, Moos MK, et al. Measuring the intensity of pregnancy planning effort. *Paediatr Perinat Epidemiol* 2003;**17**:97–105.
- Cartwright A. *Parents and family planning services*. London: Routledge Kegan Paul, 1970.
- Bone M. *Family planning services in England and Wales*. London: HMSO, 1973.
- Dunnell K. *Family formation 1976*. London: HMSO, 1979.
- Fleissig A. Unintended pregnancies and the use of contraception: changes from 1984 to 1989. *BMJ* 1991;**302**:147.
- Cleland J, Johnson-Acsadi G, Marckwardt A. The core questions. In: Cleland J, Scott C, eds. *The World Fertility Survey: an assessment*. Oxford: Oxford University Press, 1987.
- Macro International. *Women's lives and experiences: a decade of research findings from the Demographic and Health Surveys*. Program Calverton, MD: Macro International, 1994.
- London K, Peterson L, Piccinino L. The National Survey of Family Growth: principal source of statistics on unintended pregnancy: supplement to chapter two. In: Brown SS, Eisenberg L, eds. *The best intentions: unintended pregnancy and the well-being of children and families*. Washington: National Academy Press, 1995.
- Adams MM, Shulman HB, Bruce C, et al. The Pregnancy Risk Assessment Monitoring System: design, questionnaire, data collection and response rates. *Paediatr Perinat Epidemiol* 1991;**5**:333–46.
- Kost K, Forrest JD. Intention status of U.S. births in 1988: differences by mothers' socioeconomic and demographic characteristics. *Fam Plan Perspect* 1995;**27**:11–17.
- Sable MR, Spencer JC, Stockbauer JW, et al. Pregnancy wantedness and adverse pregnancy outcomes: differences by race and Medicaid status. *Fam Plan Perspect* 1997;**29**:76–81.
- Kaufmann RB, Morris L, Spitz AM. Comparison of two question sequences for assessing pregnancy intentions. *Am J Epidemiol* 1997;**145**:810–16.
- Trussell J, Vaughan B, Stanford J. Are all contraceptive failures unintended pregnancies? Evidence from the 1995 National Survey of Family Growth. *Fam Plan Perspect* 1999;**31**:246–7, 260.
- Stanford JB, Hobbs R, Jameson P, et al. Defining dimensions of pregnancy intendedness. *Maternal Child Health Journal* 2000;**4**:183–9.
- Petersen R, Moos MK. Defining and measuring unintended pregnancy: issues and concerns. *Womens Health Issues* 1997;**7**:234–40.
- Luker K. Contraceptive failure and unintended pregnancy: a reminder that human behavior frequently refuses to conform to models created by researchers. *Fam Plan Perspect* 1999;**31**:248–9.
- Santelli J, Rochat R, Hatfield-Timajchy K, et al. The measurement and meaning of unintended pregnancy. *Perspect Sex Repro Health* 2003;**35**:94–101.
- Peterson LS, Mosher WD. Contraceptive failure and unintended pregnancy: options for measuring unintended pregnancy in Cycle 6 of the National Survey of Family Growth. *Fam Plan Perspect* 1999;**31**:252–3.
- Barrett G. *Developing a measure of unplanned pregnancy*. [PhD thesis]. London: University of London, 2002.
- Barrett G, Wellings K. What is a "planned" pregnancy? Empirical data from a British study. *Soc Sci Med* 2002;**55**:545–57.
- Everett CB. Unintended pregnancies and contraceptive use. *BMJ* 1991;**302**:789–90.
- Joyce T, Kaestner R, Korenman S. On the validity of retrospective assessments of pregnancy intention. *Demography* 2002;**39**:199–213.
- Nunnally JC, Bernstein IH. *Psychometric theory*. 3rd edn. New York: McGraw-Hill, 1994.
- Cronbach LJ. Coefficient alpha and the internal structure of tests. *Psychometrika* 1951;**16**:297–334.
- Loewenthal KM. *An introduction to psychological tests and scales*. 2nd edn. London: Psychology Press, 2001.
- Kline P. *A handbook of test construction: introduction to psychometric design*. London: Methuen, 1986.
- Streiner DL, Norman GR. *Health measurement scales: a practical guide to their development and use*. 2nd edn. Oxford: Oxford University Press, 1995.
- DeVellis RF. *Scale development: theory and applications*. London: Sage, 1991.
- Ware JE, Snow KK, Kosinski M, et al. *SF-36 Health Survey: manual and interpretation guide*. Boston: The Health Institute, 1993.
- Kincaid JP, Fishburne RP, Rogers RL, et al. *Derivation of new readability formulas (Automated Readability Index, Fog Count, and Flesch Reading Ease Formula) for navy enlisted personnel*. Research Branch report 8–75. Memphis: Naval Air Station, 1975.
- Scientific Advisory Committee of the Medical Outcomes Trust. Assessing health status and quality-of-life instruments: attributes and review criteria. *Qual Life Res* 2002;**11**:193–205.
- Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;**33**:159–74.
- Kline P. *The new psychometrics: science, psychology and measurement*. London: Routledge, 1998.
- Ware JE, Keller SD. Interpreting general health measures. In: Spilker B, ed. *Quality of life and pharmacoeconomics*. 2nd edn. Philadelphia: Lippincott-Raven Publishers, 1996.
- Office for National Statistics. *Birth statistics: review of the registrar general on births and patterns of family building in England and Wales, 1999*. Series FM1 no 28. London: HMSO, 2000.
- Office for National Statistics. *Abortion statistics: legal abortions carried out under the 1967 Abortion Act in England and Wales, 1999*. Series AB, no 26. London: HMSO, 2000.
- General Register Office for Scotland. Annual Report, 1999. (<http://www.gro-scotland.gov>).
- Information and Statistics Division of NHS in Scotland. *Abortion Statistics Scotland 1998 and 1999*. Health Briefing, number 00/11, July, 2000.
- Streiner DL. Sample-size formulae for parameter estimation. *Percept Mot Skills* 1994;**78**:275–84.
- Donner A, Eliasziw M. Sample size requirements for reliability studies. *Stat Med* 1987;**6**:441–8.

- 45 **Office for National Statistics.** *Birth statistics: review of the registrar general on births and patterns of family building in England and Wales, 2000*, Series FM1 no 29. London: HMSO, 2001.
- 46 **Office for National Statistics.** *Abortion statistics: legal abortions carried out under the 1967 Abortion Act in England and Wales, 2000*. Series AB, no 27. London: HMSO, 2001.
- 47 **General Register Office for Scotland.** Annual report, 2000. (<http://www.gro-scotland.gov>).
- 48 **Szreter S.** *Fertility, class and gender in Britain, 1860–1940*. Cambridge: Cambridge University Press, 1996.
- 49 **Fisher K.** Uncertain aims and tacit negotiation: birth control practices in Britain, 1925–50. *Popul Dev Rev* 2000;**26**:295–317.

THE JECH GALLERY

doi: 10.1136/jech.2003.008730

Influential women in occupational health

Lynn R Goldman, MD—Paediatrician, Professor, and Epidemiologist



April 1951–, Country of birth: USA

Lynn Goldman has extensive background in pesticide and environmental risks to children. During her tenure as Assistant Administrator for Prevention, Pesticides and Toxic Substances at the US Environmental Protection Agency (EPA), she was responsible for the nation's toxic chemicals regulatory programmes. There she expanded citizens' right to know under the Toxics Release Inventory, and updated the US pesticides laws. Goldman made progress on testing of high volume industrial chemicals and identification of chemicals that disrupt endocrine systems. She was also successful in promoting children's health issues, furthering the international agenda for global chemical safety.

“The environment has been taken out of public health and put into environmental agencies, and environmental agencies have lost touch with health.”

With globalisation, Goldman recognises the increasing importance of chemical safety and biotechnology to public health. Her work involves development of policy. At Johns Hopkins University, she co-chairs the school-wide taskforce on bioterrorism.

D F Salerno

Clinical Communications Scientist, Pfizer Global Research and Development, Ann Arbor Laboratories, Ann Arbor, MI, USA

I L Feitshans

Adjunct Faculty, Cornell University, School of Industrial and Labor Relations, Albany, NY, USA

Correspondence to: Deborah F. Salerno, 2800 Plymouth Road, Ann Arbor, MI 48105, USA; deborah.salerno@pfizer.com